

ISCR Inhibitor Screen

Synonyms

Clinical Indication

Used to establish the presence of immediate acting or time-dependent clotting factor specific inhibitor.

Clinically indicated from prolonged APTT and / or prolonged PT

Part of Profile / See Also

Request Form

Combined Pathology manual Blood form or ICE request

Availability / Frequency of Analysis

On request after discussion with consultant haematologist.

Referred test: Analysed at Royal London ([Barts Health 8285](#)) if specific criteria met.

Turnaround Time

7 days

Patient Preparation

DO NOT use heparinised syringes / lines for collecting coagulation screens. Patient should be bled with minimal stasis, and samples received in lab within 2 hours of collection.

Sample Requirements

Specimen Type

Citrated whole blood

Volume

Collection tube must be filled to 'fill mark' on side of bottle. This is critical.

Container



2 x Blue top (sodium citrate) tube



2 x Paediatric Blue Top (sodium citrate) tubes

Samples must be received by lab within 2 hours of collection.

Reference Range & Units

See test reports

Interferences

Activated sample, sample haemolysis or lipaemia.

Interpretation & Clinical

See test reports

Decision Value (if applicable)

References

Test code

RLTA

Lab Handling

Samples to be double spun and frozen.



**Accredited to
ISO 15189:2022**