



ISCR Inhibitor Screen

Synonyms	
Clinical Indication	<p>Used to establish the presence of immediate acting or time-dependent clotting factor specific inhibitor.</p> <p>Clinically indicated from prolonged APTT and / or prolonged PT</p>
Part of Profile / See Also	
Request Form	Combined Pathology manual Blood form or ICE request
Availability / Frequency of Analysis	<p>On request after discussion with consultant haematologist.</p> <p>Referred test: Analysed at Royal London (Barts Health 8285) if specific criteria met.</p>
Turnaround Time	7 days
Patient Preparation	DO NOT use heparinised syringes / lines for collecting coagulation screens. Patient should be bled with minimal stasis, and samples received in lab within 2 hours of collection.
Sample Requirements	
Specimen Type	Citrated whole blood
Volume	Collection tube must be filled to 'fill mark' on side of bottle. This is critical.
Container	 2 x Blue top (sodium citrate) tube  2 x Paediatric Blue Top (sodium citrate) tubes <p>Samples must be received by lab within 2 hours of collection.</p>
Reference Range & Units	See test reports
Interferences	Activated sample, sample haemolysis or lipaemia.
Interpretation & Clinical Decision Value (if applicable)	See test reports
References	
Test code	RLTA
Lab Handling	Samples to be double spun and frozen.