



PF-PTD-194

Joint Fluid

Synonyms	Joint aspirate
Clinical Indication	Septic arthritis
	Please give detailed clinical information, including date of onset and any current or intended antibiotic therapy
Part of Profile / See Also	Microscopy for organisms & WBCs WBC count for Southend Trust only (not performed if sample clotted or heavily blood stained). WBC semi quantitative (i.e., +/-, +, +++, +++) estimation for BTUH samples or if sample clotted or heavily blood stained WBC differential if cells are ≥ ++ or ≥ 50 wbc/cmm Requests for crystals will be referred to Cytology (a separate sample is desirable to be sent directly to Cytology) Crystals will be performed on all Basildon Trust samples. Crystals will be by request only on Southend Trust samples or if seen on microscopy. Culture for aerobic and anaerobic organisms.
Request Form	Order comms request i.e., ICE or Medway or clearly handwritten Microbiology request forms where order comms not available
Availability / Frequency of	During working day or out of hours as specified below.
Analysis	For urgent requests, please contact laboratory 01268 968285
	Out of hours, please contact on call Microbiology lab staff via switchboard
	Basildon requests will only be accepted up to 22:00hrs.
Turnaround Time	Microscopy: Same day.
. Gridiodila IIIIo	
	Culture: 5 days For enrichment of samples i.e., where blood culture bottles are submitted, these joint fluids are incubated for 14 days followed by terminal sub therefore may be subject to a prolonged TAT for a final negative report.
Patient Preparation	
Sample Requirements	Refrigerate sample in universal if there is to be a delay in delivery to the
	laboratory.
	Blood culture bottles NOT to be refrigerated- bottles can be loaded onto the
	FX at the ESL.
Specimen Type	Joint Fluid
Volume	20 ml - Minimum of 1 ml.
	Blood culture bottle - fill according to volume specified on bottle





PF-PTD-194

Container

White capped sterile universal - CE marked



Blood culture bottle sets inoculated directly at point of collection may be submitted in addition to the universal container.



Reference Range & Units

Interferences

Interpretation & Clinical

Decision Value (if applicable)

Positive culture results will be reported with antibiotic susceptibilities where appropriate.

Microscopy results will be phoned to the requestor out of hours.

References

Test code

JOIF

Lab Handling

Processed on receipt.