


## Antibody Titre / Quantitation

<b>Synonyms</b>	
<b>Clinical Indication</b>	As per BCSH Guidelines for Blood Grouping and Antibody Testing During Pregnancy. Antibody must be present.
<b>Part of Profile / See Also</b>	
<b>Request Form</b>	Combined Pathology manual Blood form, Blood Transfusion form or ICE request
<b>Availability / Frequency of Analysis</b>	Analysed by referral laboratory, NBS Colindale, if specific criteria met.
<b>Turnaround Time</b>	3-7 days
<b>Patient Preparation</b>	
<b>Sample Requirements</b>	
<b>Specimen Type</b>	Plasma / Whole Blood
<b>Volume</b>	3 x 6ml EDTA Sample
<b>Container:</b>	
Basildon Site:	 Purple top (EDTA) tube.
Southend Site:	 Pink Top (EDTA) tube.
	<b>Samples should be transported to laboratory immediately.</b>
<b>Reference Range &amp; Units</b>	
<b>Interferences</b>	Insufficient sample quantity. Sub detectable antibody.
<b>Interpretation &amp; Clinical Decision Value (if applicable)</b>	As per BCSH Guidelines for Blood Grouping and Antibody Testing During Pregnancy
<b>References</b>	
<b>Test code</b>	
<b>Lab Handling</b>	Samples should be transported as soon as possible